

$\pi\pi$ scattering in a nonlocal Nambu – Jona-Lasinio model

A. A. Osipov,^{1,*} A. E. Radzhabov,^{2,†} and M. K. Volkov^{2,‡}

¹*Dzhelepov Laboratory of Nuclear Problems,*

Joint Institute for Nuclear Research, 141980 Dubna, Russia

²*Bogoliubov Laboratory of Theoretical Physics,*

Joint Institute for Nuclear Research, 141980 Dubna, Russia

We consider a nonlocal version of the Nambu and Jona-Lasinio model. The non-locality is contained in the quark-antiquark bilinears of the four-quark vertices as a form factor of the Gaussian type. The model has three parameters which can be fixed in favour of the values of the pion mass, the pion decay constant f_π , and the current quark mass. The $\pi\pi$ scattering amplitude is obtained by calculating the quark box and the σ -pole diagrams, where σ is the scalar isoscalar meson. It is shown that this amplitude satisfies the well-known Weinberg relation. We obtain the s, p, d wave scattering lengths in all isotopic channels and the s wave slope parameters. The results are in satisfactory agreement with both phenomenological data and the basic requirements of low-energy theorems, thus supporting to the form factor used.

1. INTRODUCTION

The theoretical study of $\pi\pi$ scattering has a long-standing history. One of the pioneering works which has certainly boosted the development of chiral symmetric models was due to Weinberg [1]. In that paper the methods of current algebra [2, 3] were successfully used to compute the low-energy $\pi\pi$ scattering lengths. After that many Lagrangian models of strong interactions of hadrons were intensively developed, using the linear [4] and nonlinear [5–7] realizations of chiral symmetry. Various low-energy properties of hadrons were described in the framework of these models, for instance $\pi\pi$ scattering (see e.g. [8–10]).

*Electronic address: osipov@nu.jinr.ru

†Electronic address: aradzh@theor.jinr.ru

‡Electronic address: volkov@theor.jinr.ru

The chiral perturbation theory has been developed on the basis of nonlinear Lagrangians. This effective field theory describes the low-energy structure of different amplitudes in terms of expansion in powers of energies, momenta and current quark masses [11, 12]. Chiral symmetry determines the low energy behaviour of the $\pi\pi$ scattering amplitude to within very small uncertainties [13, 14]. For such a theory one takes for granted that chiral symmetry is spontaneously broken. There are theoretical arguments indicating that chromodynamics leads indeed to the formation of a quark condensate [15]. However, the chiral perturbation theory cannot explain this phenomenon.

An analogy with superconductivity led Nambu and Jona-Lasinio (NJL) to the field theoretical model of strong four-fermion interactions [16, 17] which provide for spontaneous chiral symmetry breaking and the formation of a quark condensate. The local version of this model was particularly well developed at the quark level [18–25]. The spectrum, low-energy dynamics, the main strong and electromagnetic decays, scattering and the internal characteristics of mesons have a reasonable explanation within this model.

In this paper, we generalize the local approach to $\pi\pi$ scattering by constructing the $\pi\pi$ scattering amplitude within the framework of a nonlocal NJL model with linear realization of chiral symmetry. The scalar and pseudoscalar degrees of freedom are considered. Let us note that one needs to know higher powers of the p^2 expansion to describe scattering lengths and slope parameters of d -waves and higher. Such expansions were first a problem in the local approach: ad hoc form factors were inserted into the meson-quark vertices [26], their structure not being derived from the internal structure of the model, thus requiring justification [27].

A coherent method which takes into account the whole contribution of quark one-loop diagrams in the local NJL model has been formulated in [28, 29]. In particular, the method was successfully used to calculate $\pi\pi$ scattering lengths.

Recently, nonlocal versions of the NJL model have been actively developed [30–42]. These models are free from ultra-violet divergences, as well as there is hope to confine the quarks because the constituent quark mass depends on the momentum. The form factors of the meson-quark vertices are the direct consequence of this nonlocality. Thus, it seems interesting to calculate again $\pi\pi$ scattering lengths and parameters of effective ranges in the framework of a nonlocal model. The relative simplicity of the process together with a large amount of experimental information about $\pi\pi$ scattering serve us to gain a considerable

insight into the structure of nonlocal interactions from such calculations.

The paper is organized as follows. In the next Sect. we consider the effective $SU(2) \times SU(2)$ symmetric nonlocal four-quark interaction. The dynamical mechanism of spontaneous chiral symmetry breaking is clarified after bosonization. We show that the model satisfies Goldberger – Treiman and Gell-Mann, Oaks, Renner relations. We fix parameters of the model in this section. In Sect. 3 we obtain the $\pi\pi$ scattering amplitude, check Weinberg relation and calculate s , p , d scattering lengths in all isotopic channels and slope parameters of s -waves. The results are compared with empirical data. In the last Sect. we discuss them.

2. NONLOCAL MODEL OF THE NJL TYPE

The model is based on the effective $SU(2) \times SU(2)$ symmetric quark interaction in the scalar and pseudoscalar sectors

$$\mathcal{L} = \bar{q}(x)(i\hat{\partial}_x - m^0)q(x) + \frac{G}{2} (J_\sigma(x)J_\sigma(x) + J_\pi^a(x)J_\pi^a(x)), \quad (1)$$

where $q, \bar{q}$ are quark and antiquark fields, m are masses of current quarks, τ_i are the Pauli matrices and G is the four-quark coupling constant. The nonlocal quark currents $J_I(x)$ are given by¹

$$J_I(x) = \int d^4x_1 d^4x_2 \delta\left(x - \frac{x_1 + x_2}{2}\right) f((x_1 - x_2)^2) \bar{q}(x_1) \Gamma_I q(x_2), \quad (2)$$

where the function $f(x)$ is normalized as $f(0) = 1$. The matrices Γ_I in eq.(1) are defined as follows $\Gamma_\sigma = \mathbf{1}$, $\Gamma_\pi^a = i\gamma^5 \tau^a$.

One has for this Lagrangian after bosonization

$$\mathcal{L} = \bar{q}(x)(i\hat{\partial}_x - m_c)q(x) - \frac{1}{2G} (\pi^a(x)^2 + \tilde{\sigma}(x)^2) + J_\sigma(x)\tilde{\sigma}(x) + \pi^a(x)J_\pi^a(x), \quad (3)$$

where $\tilde{\sigma}$ and π^a are the scalar and pseudoscalar fields. The scalar field $\tilde{\sigma}$ has a nonzero vacuum expectation value $\langle \tilde{\sigma} \rangle_0 = \sigma_0 \neq 0$. To arrive at the physical scalar field with zero expectation value, it is necessary to shift the field: $\tilde{\sigma} = \sigma + \sigma_0$.

Varying the action with respect to σ

$$\left\langle \frac{\delta \mathcal{S}}{\delta \sigma} \right\rangle_0 = 0, \quad (4)$$

¹ The models with such nonlocality have been considered previously in [37–42]

one obtains the equation for the dynamical quark mass, the “gap” equation

$$m(p^2) = m_c + iG \frac{N_f N_c}{(2\pi)^4} f(p^2) \int \frac{d^4 k}{(2\pi)^4} f(k^2) \text{Tr} [S(k)], \quad (5)$$

where N_f and N_c are the numbers of quark flavours and colours, respectively. Note that due to the form factors the quark loop integral is finite. This equation can also be written as

$$m(p^2) = m_c - \sigma_0 f(p^2) = m_c + (m_q - m_c) f(p^2), \quad (6)$$

where m_q is a dimensionfull parameter which plays the role of the constituent quark mass. As a result the quark mass becomes p^2 dependent. The quark propagator $S(p)$ has the form

$$S(p) = \frac{1}{\hat{p} - m(p^2)}. \quad (7)$$

The propagators of the meson fields are

$$D_{\sigma,\pi}(p^2) = \frac{1}{-G^{-1} + J_{\sigma,\pi}(p^2)} = \frac{g_{\sigma,\pi}^2(p^2)}{p^2 - M_{\sigma,\pi}^2}, \quad (8)$$

where $M_{\sigma,\pi}$ are meson masses, $g_{\sigma,\pi}(p^2)$ are the functions describing the renormalization of mesonic fields, and $J_{\sigma,\pi}(p^2)$ are the contributions of the quark one-loop diagrams

$$J_{\sigma,\pi}(p^2) = i \frac{N_f N_c}{(2\pi)^4} \int d^4 k f^2(k^2) \text{Tr} [S(k_-) \Gamma_{\sigma,\pi} S(k_+) \Gamma_{\sigma,\pi}], \quad (9)$$

where $k_+ = k + p/2$, $k_- = k - p/2$. The meson masses $M_{\sigma,\pi}$ are the poles of the propagators

$$J_{\sigma,\pi}(M_{\sigma,\pi}^2) = G^{-1}, \quad (10)$$

and the couplings $g_{\sigma,\pi}(M_{\sigma,\pi}^2)$ are determined on the mass-shell of the mesonic states by the formulae

$$g_{\sigma,\pi}^{-2}(M_{\sigma,\pi}^2) = \left. \frac{dJ_{\sigma,\pi}(p^2)}{dp^2} \right|_{p^2=M_{\sigma,\pi}^2}. \quad (11)$$

The amplitude of the pion decay $\pi \rightarrow \mu\nu$ is

$$A_{(\pi \rightarrow \mu\nu)}^\mu(p) = ip^\mu f_\pi, \quad (12)$$

where f_π is the coupling of the weak pion decay². The Goldberger – Treiman relation is held in the chiral limit $f_\pi = m_q/g_\pi(0)$ and f_π is given by the formula of Pagels and Stokar [45]

$$f_\pi^2 = \frac{N_c}{4\pi^2} \int_0^\infty du u m(u) \frac{m(u) - \frac{1}{2} u m'(u)}{(u + m^2(u))^2}. \quad (13)$$

² Detailed description of insertion of electroweak fields into nonlocal Lagrangian is given in appendix.




Figure 1. Diagrams contributing to the amplitude of $\pi\pi$ scattering.

This expression is given in Euclidean domain $u = -p^2$.

The Gell-Mann – Oaks – Renner relation is fulfilled in the model

$$M_\pi^2 f_\pi^2 = -2 m_c \langle \bar{q}q \rangle + O(m_c^2), \quad (14)$$

where $\langle \bar{q}q \rangle$ is the quark condensate. This relation can be obtained from the equation for the pion mass (10) with the help of the gap equation (5) and expanding in powers of M_π^2 and m_c . The form factor $f(u)$ has been chosen in the form of a Gaussian exponent $f(u) = \exp(-u/\Lambda^2)$.

The model has three parameters m_c, Λ, G . After using the values of pion mass $M_\pi = 140$ MeV and the weak pion decay constant $f_\pi = 92$ MeV as input parameters we obtain instead of three arbitrary parameters only one m_c . This parameter can be chosen in the interval $4.05 - 7.425$ MeV at the typical hadron scale 1 GeV [43]. Additional restriction can be obtained, e.g., from the d -wave scattering length a_2^2 . As a result, the parameters of the model are $m_c = 4.8$ MeV, $\Lambda = 880$ MeV and $G = 22.87$ GeV $^{-2}$. Thus, one can find that $m_q = 345$ MeV and $\langle \bar{q}q \rangle = (-255 \text{ MeV})^3$. For the mass and the width of scalar meson we obtain $M_\sigma = 570$ MeV, $\Gamma_\sigma = 220$ MeV.

3. $\pi\pi$ SCATTERING

The amplitude of the process is described by two main diagrams plotted on Fig.1.

Using the notation of Chew and Mandelstam one can write the matrix element of $\pi\pi$ scattering

$$\langle i_1 i_2 | \mathcal{A} | i_3 i_4 \rangle = \delta_{i_1 i_2} \delta_{i_3 i_4} A(s, t, u) + \delta_{i_1 i_3} \delta_{i_2 i_4} A(t, s, u) + \delta_{i_1 i_4} \delta_{i_2 i_3} A(u, t, s) \quad (15)$$

in terms of three kinematical invariants

$$s = (p_1 + p_2)^2, \quad t = (p_1 - p_3)^2, \quad u = (p_1 - p_4)^2. \quad (16)$$

For the amplitudes with the definite isospin values $I = 0, 1, 2$ one has

$$\begin{aligned} T^0 &= 3A(s, t, u) + A(t, s, u) + A(u, t, s) \\ T^1 &= A(t, s, u) - A(u, t, s) \\ T^2 &= A(t, s, u) + A(u, t, s). \end{aligned} \tag{17}$$

The σ -pole diagram contains the $\sigma\pi\pi$ vertex which has the following structure:

$$\begin{aligned} A_{\sigma \rightarrow \pi\pi} &= 2g_\sigma(p^2)g_\pi(p_1^2)g_\pi(p_2^2)g_{\sigma\pi\pi}(p^2, p_1^2, p_2^2) \\ g_{\sigma\pi\pi}(p^2, p_1^2, p_2^2) &= i\frac{N_f N_c}{(2\pi)^4} \int d^4k f\left(\left(k + \frac{p_1}{2}\right)^2\right) f\left(\left(k - \frac{p_2}{2}\right)^2\right) f\left(\left(k + \frac{p_1 - p_2}{2}\right)^2\right) \\ &\quad \times \text{Tr}[S(k + p_1)\Gamma_\sigma S(k - p_2)\Gamma_\pi S(k)\Gamma_\pi], \end{aligned}$$

The box diagram contributes as

$$\begin{aligned} A_{box}(s, t, u) &= 2g_\pi^4(M_\pi^2)(J_4(s, t, u) + J_4(s, u, t) - J_4(u, t, s)) \\ J_4(s, t, u) &= i\frac{N_f N_c}{(2\pi)^4} \int d^4k f\left(\left(k + \frac{p_1}{2}\right)^2\right) f\left(\left(k - \frac{p_2}{2}\right)^2\right) f\left(\left(k + \frac{p_1 - p_3}{2}\right)^2\right) \\ &\quad \times f\left(\left(k + \frac{p_1 - p_2 - p_3}{2}\right)^2\right) \text{Tr}[S(k)\Gamma_\pi S(k - p_2)\Gamma_\pi S(k + p_1 - p_3)\Gamma_\pi S(k + p_1)\Gamma_\pi], \end{aligned}$$

As a result, the whole $\pi\pi$ amplitude can be written as

$$A(s, t, u) = A_{box}(s, t, u) + 4g_\pi^4(M_\pi^2)\frac{g_{\sigma\pi\pi}^2(s, M_\pi^2, M_\pi^2)}{G^{-1} - J_\sigma(s)}. \tag{18}$$

At the lowest order of the momentum and m_c expansions one obtains the Weinberg result

$$A(s, t, u) = \frac{s - M_\pi^2}{f_\pi^2} + O(p^4) \tag{19}$$

Let us recall that in the local NJL model the Weinberg formula can be obtained on the basis of the box and the σ -pole diagrams, if one picks up only the divergent terms of the quark one-loop diagrams [26, 46]

$$A(s, t, u) = -4g^2 + \frac{16m^2g^2}{M_\sigma^2 - s} = \frac{s - M_\pi^2}{f_\pi^2} + \dots, \tag{20}$$

where $g = g_\pi = g_\sigma$, m is the constituent quark mass, and the sigma meson mass takes its model value $M_\sigma^2 = M_\pi^2 + 4m^2$.

Scattering lengths a_l^I and slope parameters b_l^I are the first terms in the expansion of partial-wave amplitudes for orbital angular momentum l and isospin I

$$\lim_{k^2 \rightarrow 0} \frac{1}{64\pi M_\pi k^{2l}} \int_{-1}^1 dx P_l(x) T^I(s, t, u) = a_l^I + k^2 b_l^I + O(k^4). \quad (21)$$

Here $s = 4(M_\pi^2 + k^2)$, $t = -2k^2(1 - x)$, $u = -2k^2(1 + x)$; k is the c.m. three-momentum, and $x = \cos \theta$, where θ is a scattering angle between pions in the c.m. frame and $P_l(x)$ are the Legendre polynomials.

To obtain scattering lengths we expand the integrand of eq.(21) in powers of k^2 . The subsequent integration over x lead to the expressions for the scattering lengths and slope parameters that are given in appendix.

The obtained values for the scattering lengths of the s, p and d waves in all isotopic channels as well as the parameters of the effective ranges for the s waves are given in Table. To compare our results, we also present there characteristics obtained in the framework of the local NJL model [28]. The empirical data are taken from the papers [49] (the life-time of the $\pi^+\pi^-$ mesonic atom), [50] (the $K \rightarrow 3\pi$ decay), [48] (the $K^\pm \rightarrow \pi^+\pi^- e^\pm \nu_e(\bar{\nu}_e)$ decay), and [47] (the phenomenological analysis of data).

To check the result, we passed to the local limit in our formulae, by regularizing the quark one-loop diagrams in the standard way and taking $\Lambda \rightarrow \infty$ in the nonlocal form factors. After this the results of known calculations in the local model are reproduced [28].

4. CONCLUSIONS

The present calculations show that the main low energy theorems of pion physics are fulfilled within the nonlocal NJL model extension considered. We have checked the Goldberger – Treiman relation, the Gell-Mann, Oaks, Renner formula and the famous Weinberg result for the low energy $\pi\pi$ scattering amplitude.

The Goldberger – Treiman relation is an exact result of the theory in the Goldstone limit and reflects the chiral asymmetry of the ground state. In the NJL model as well as in the linear sigma model, the Goldberger – Treiman relation can be understood as $\langle \sigma_0 \rangle = f_\pi$, where $\langle \sigma_0 \rangle$ represents the point at which the effective potential has a minimum.

The Gell-Mann, Oaks, Renner formula in the NJL model relates the quark condensate with the gap equation, thus reflecting the dynamical mechanism for the spontaneous chiral

Table 1. Scattering lengths and slope parameters for the $\pi\pi$ system. Columns LNJL, PY and EXP correspond to the results obtained in the local NJL model [28], results of the phenomenological analysis of experimental data [47], and known data extracted from the life-time of $\pi^+\pi^-$ mesonic atom [49], from the $K \rightarrow 3\pi$ decay [50] and $K^\pm \rightarrow \pi^+\pi^-e^\pm\nu_e(\bar{\nu}_e)$ data [48].

	This work	LNJL	PY	EXP
$a_0^0 M_\pi$	0.20	0.190	0.230 ± 0.015	0.216 ± 0.017 [48]
$a_0^2 M_\pi$	-0.049	-0.044	-0.048 ± 0.0046	-0.0454 ± 0.0049 [48] -0.041 ± 0.036 [50]
$(a_0^0 - a_0^2) M_\pi$	0.25	0.234	0.277 ± 0.014	$0.264^{+0.033}_{-0.020}$ [49] 0.268 ± 0.027 [50]
$a_1^1 \times 10^{-3} M_\pi^3$	37	34	38.4 ± 0.8	
$a_2^0 \times 10^{-4} M_\pi^5$	16.3	16.7	18.7 ± 0.41	
$a_2^2 \times 10^{-4} M_\pi^5$	2.8	3.2	2.78 ± 0.37	
$b_0^0 M_\pi^3$	0.26	0.27	0.312 ± 0.014	
$b_0^2 M_\pi^3$	-0.085	-0.079	-0.090 ± 0.006	

symmetry breaking.

The $\pi\pi$ scattering amplitude has been obtained in the Hartree-Fock approximation of the nonlocal NJL model and was used to calculate the threshold characteristics. The results for the scattering lengths in s, p and d waves and for the s -wave slope parameters are in satisfactory agreement with the empirical data.

The main contribution to the values of the s, p scattering lengths is given by the p^2 terms of the amplitudes. By virtue of the Weinberg relation, they are less sensitive to the structure of the form factor. The d -waves are mainly determined by the p^4 order terms of the amplitude and the sufficient predictions for the d -waves are a non-trivial result of our calculations. In terms of box and σ -pole diagrams, they are additive for the a_2^0 scattering length and have opposite signs in the a_2^2 case. The value of the a_2^2 scattering lengths is strongly affected by the magnitude of the σ meson mass predicted by the model.

Returning to the problem of fixing model parameters, note that the best agreement with experimental a_2^2 is achieved for $m_c = 4.8$ MeV ($a_2^2 = 2.8 \times 10^{-4} M_\pi^{-5}$). For the current quark mass equal to 4.4 and 5.2 MeV this scattering length equals $3.5 \times 10^{-4} M_\pi^{-5}$ and $2 \times 10^{-4} M_\pi^{-5}$,

respectively.

In the future we plan to extend the nonlocal model considered by including the vector and axial-vector mesons and taking into account the $\pi - a_1$ transitions. It also will be interesting to consider the $U(3) \times U(3)$ chiral symmetry group and study the internal structure and interactions of the pseudoscalar, scalar, vector, and axial-vector nonets. Of special interest will be processes in which the p^2 dependence of the amplitudes will be relevant.

Moreover, it is interesting to investigate the $\pi\pi$ scattering process in the hot and dense medium in the framework of a nonlocal model. When chiral symmetry is restored, the σ meson mass tends to the mass of pion [42, 46, 51]. This can lead to a noticeable increase in the $\pi\pi$ scattering amplitude near the phase transition of hadron matter into quark-gluon plasma. Such investigations in the framework of local models are performed in [52].

The authors thank A. E. Dorokhov for useful discussions.

A.E.R. and M.K.V. acknowledge the support of the Russian Foundation for Basic Research, under contract 05-02-16699.

A. A. Osipov acknowledges support from grants provided by Fundação para a Ciência e a Tecnologia, POCTI/FNU/50336/2003 and POCI/FP/63412/2005. This research is part of the EU integrated infrastructure initiative Hadron Physics project under contract No.RII3-CT-2004-506078.

APPENDIX

Weak pion decay constant

To determine the weak pion decay constant one inserts external sources associated with the electroweak fields. For this purpose it is necessary to take into account the contributions induced by the nonlocal interaction. We use the standard procedure: the external fields are introduced via the Schwinger phase factor which delocalizes quark fields

$$q(y) \rightarrow Q(x, y) = \mathcal{P} \exp \left\{ i \int_x^y dz^\mu [V_\mu^a(z) + A_\mu^a(z) \gamma_5] T^a \right\} q(y). \quad (22)$$

Here $\mathcal{P}$ is the Dyson ordering along the path, $V_\mu^a(z)$ and $A_\mu^a(z)$ are the external vector and axial-vector gauge fields which can be identified with electroweak fields, T^a stands for the generators of the flavour group. Since the derivative of the path integral of any function




Figure 2. Diagrams corresponding to the weak decay of the pion $\pi \rightarrow \mu\nu$.

$F_\mu(z)$ over the path does not depend on the trajectory along which one calculates the integral [44]

$$\frac{\partial}{\partial y^\mu} \int_x^y dz^\nu F_\nu(z) = F_\mu(y), \quad \delta^{(4)}(x-y) \int_x^y dz^\nu F_\nu(z) = 0. \quad (23)$$

As a result, non-minimal interactions, which could be induced by the kinetic term of quark fields, are absent. The gauge invariant procedure used here leads to additional vertices with any number of external fields which have their origin in the nonlocal nature of the interaction.

Weak pion decay is described by three diagrams (see fig.2). Note that in the chiral limit the contributions of nonlocal diagrams are exactly cancelled.

Scattering lengths and slope parameters

$$\begin{aligned} a_0^2 &= \frac{1}{32\pi M_\pi} \left(A(0, 4M_\pi^2, 0) + A(0, 0, 4M_\pi^2) \right) \\ (a_0^0 - a_0^2) &= \frac{3}{32\pi M_\pi} A(4M_\pi^2, 0, 0) \\ a_1^1 &= \frac{1}{48\pi M_\pi} \left(A^{(s)}(0, 4M_\pi^2, 0) + A^{(s)}(0, 0, 4M_\pi^2) - A^{(t)}(0, 0, 4M_\pi^2) - A^{(u)}(0, 4M_\pi^2, 0) \right) \\ a_2^2 &= \frac{1}{120\pi M_\pi} \left(A^{(u,u)}(0, 4M_\pi^2, 0) + A^{(t,t)}(0, 0, 4M_\pi^2) - 2A^{(s,u)}(0, 4M_\pi^2, 0) - \right. \\ &\quad \left. - 2A^{(s,t)}(0, 0, 4M_\pi^2) + A^{(s,s)}(0, 0, 4M_\pi^2) + A^{(s,s)}(0, 4M_\pi^2, 0) \right) \\ (a_2^0 - a_2^2) &= \frac{1}{40\pi M_\pi} \left(A^{(u,u)}(4M_\pi^2, 0, 0) - 2A^{(t,u)}(4M_\pi^2, 0, 0) + A^{(t,t)}(4M_\pi^2, 0, 0) \right) \end{aligned}$$

$$b_0^2 = \frac{1}{16\pi M_\pi} \left(2A^{(u)}(0, 0, 4M_\pi^2) + 2A^{(t)}(0, 4M_\pi^2, 0) - A^{(u)}(0, 4M_\pi^2, 0) \right. \\ \left. - A^{(t)}(0, 0, 4M_\pi^2) - A^{(s)}(0, 4M_\pi^2, 0) - A^{(s)}(0, 0, 4M_\pi^2) \right) \\ (b_0^0 - b_0^2) = \frac{3}{16\pi M_\pi} \left(2A^{(s)}(4M_\pi^2, 0, 0) - A^{(t)}(4M_\pi^2, 0, 0) - A^{(u)}(4M_\pi^2, 0, 0) \right),$$

where

$$A^{(x)}(a, b, c) = \frac{\partial}{\partial x} A(s, t, u)|_{s=a, t=b, u=c}, \quad A^{(x,y)}(a, b, c) = \frac{\partial^2}{\partial x \partial y} A(s, t, u)|_{s=a, t=b, u=c}.$$

-
1. S. Weinberg, Phys. Rev. Lett. **17** (1966) 616.
 2. S. L. Adler and R. F. Dashen, *Current algebra* (Benjamin, New York, 1968).
 3. V. De Alfaro, S. Fubini, G. Furlan and C. Rossetti, *Currents in hadron physics* (North-Holland, Amsterdam, 1973).
 4. S. Gasiorowicz and D. A. Geffen, Rev. Mod. Phys. **41** (1969) 531.
 5. S. Coleman, J. Wess and B. Zumino, Phys. Rev. **177** (1969) 2239.
 6. C. Callan, S. Coleman, J. Wess and B. Zumino, Phys. Rev. **177** (1969) 2247.
 7. M. K. Volkov and V. N. Pervushin, *Essentially nonlinear quantum theories, dynamical symmetries, and meson physics*, Atomizdat, Moscow, 1978, 240 p.
 8. H. Lehmann, Phys. Lett. B **41** (1972) 529.
 9. M. K. Volkov and V. N. Pervushin, Sov. J. Nucl. Phys. **20** (1975) 408 [Yad. Fiz. **20** (1974) 762].
 10. M. K. Volkov and V. N. Pervushin, Nuovo Cim. A **27** (1975) 277.
 11. S. Weinberg, Physica A **96** (1979) 327.
 12. J. Gasser and H. Leutwyler, Annals Phys. **158** (1984) 142.
 13. G. Colangelo, J. Gasser and H. Leutwyler, Phys. Lett. B **488** (2000) 261.
 14. G. Colangelo, J. Gasser and H. Leutwyler, Nucl. Phys. B **603** (2001) 125.
 15. C. Vafa and E. Witten, Nucl. Phys. B **234** (1984) 173.
 16. Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345.
 17. Y. Nambu and G. Jona-Lasinio, Phys. Rev. **124** (1961) 246.
 18. T. Eguchi, Phys. Rev. D **14** (1976) 2755.

19. M. K. Volkov and D. Ebert, Sov. J. Nucl. Phys. **36** (1982) 736 [Yad. Fiz. **36** (1982) 1265].
20. D. Ebert and M. K. Volkov, Z. Phys. C **16** (1983) 205.
21. M. K. Volkov, Ann. Phys. (N.Y.) **157** (1984) 282.
22. M. K. Volkov, Sov.J.Part. and Nuclei **17** (1986) 186 [Fiz. Elem. Chastits. At. Yadra **17** 433].
23. D. Ebert and H. Reinhardt, Nucl. Phys. B **271** (1986) 188.
24. S. Klimt, M. Lutz, U. Vogl and W. Weise, Nucl. Phys. A **516** (1990) 429.
25. S. P. Klevansky, Rev. Mod. Phys. **64** (1992) 649.
26. M. K. Volkov and A. A. Osipov, Sov. J. Nucl. Phys. **39** (1984) 440 [Yad. Fiz. **39** (1984) 694].
27. M. K. Volkov, Phys. Atom. Nucl. **60** (1997) 997 [Yad. Fiz. **60** (1997) 1115].
28. V. Bernard, A. A. Osipov and U. G. Meissner, Phys. Lett. B **285** (1992) 119.
29. V. Bernard, A. H. Blin, B. Hiller, Y. P. Ivanov, A. A. Osipov and U. G. Meissner, Annals Phys. **249** (1996) 499.
30. A.A. Andrianov, and V.A. Andrianov, Int. J. Mod. Phys. A **8** (1993) 1981.
31. G.V. Efimov and M.A. Ivanov, *The Quark Confinement Model Of Hadrons* (IOP, Bristol, 1993).
32. R. S. Plant and M. C. Birse, Nucl. Phys. A **628** (1998) 607.
33. L.S. Celenza, B. Huang, H.S. Wang and C.M. Shakin, Phys. Rev. C **60** (1999) 025202.
34. I. V. Anikin, A. E. Dorokhov and L. Tomio, Phys. Part. Nucl. **31** (2000) 509 [Fiz. Elem. Chast. Atom. Yadra **31** (2000) 1023].
35. A. E. Radzhabov and M. K. Volkov, Eur. Phys. J. A **19** (2004) 139.
36. A. Scarpettini, D. Gómez Dumm and N. N. Scoccola, Phys. Rev. D **69** (2004) 114018.
37. H. Ito, W. W. Buck and F. Gross, Phys. Rev. C **45** (1992) 1918.
38. C. D. Roberts, R. T. Cahill, M. E. Sevier and N. Iannella, Phys. Rev. D **49** (1994) 125.
39. P. C. Tandy, Prog. Part. Nucl. Phys. **39** (1997) 117.
40. D. Blaschke, G. Burau, Y. L. Kalinovsky, P. Maris and P. C. Tandy, Int. J. Mod. Phys. A **16** (2001) 2267.
41. A. Faessler, T. Gutsche, M. A. Ivanov, V. E. Lyubovitskij and P. Wang, Phys. Rev. D **68** (2003) 014011.
42. D. Blaschke, Y. L. Kalinovsky, A. E. Radzhabov and M. K. Volkov, arXiv:hep-ph/0508264.
43. S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.
44. S. Mandelstam, Annals Phys. **19** (1962) 1.

- 45. H. Pagels and S. Stokar, Phys. Rev. D **20** (1979) 2947.
- 46. M. K. Volkov, A. E. Radzhabov and N. L. Russakovich, Phys. Atom. Nucl. **66**, 997 (2003) [Yad. Fiz. **66**, 1030 (2003)].
- 47. J. R. Pelaez and F. J. Yndurain, Phys. Rev. D **71** (2005) 074016.
- 48. S. Pislak *et al.*, Phys. Rev. D **67** (2003) 072004 [arXiv:hep-ex/0301040].
- 49. B. Adeva *et al.* [DIRAC Collaboration], Phys. Lett. B **619** (2005) 50 [arXiv:hep-ex/0504044].
- 50. J. R. Batley *et al.* [NA48/2 Collaboration], Phys. Lett. B **633** (2006) 173.
- 51. D. Ebert, Y.L. Kalinovsky, L. Munchow and M.K. Volkov, Int. J. Mod. Phys. A **8** (1993) 1295.
- 52. D. Jido, T. Hatsuda and T. Kunihiro, Phys. Rev. D **63** (2001) 011901.